



NAURU METEOROLOGICAL & HYDROLOGICAL SERVICES



The Early Action Rainfall Watch provides sector managers with a brief summary of recent rainfall patterns, particularly drought and the rainfall outlook for the coming months.

Issue Date: 13/11/2025

ENSO OUTLOOK

Current El Nino - Southern Oscillation Status

ENSO



La Niña Alert

The ENSO outlook is at ENSO La-Nina WATCH

Current **ENSO** status is at **La-Nina ALERT**. This means **La-Nina (dry conditions)** may form in the coming months.

Conditions are nearing La-Nina in the tropical Pacific, and have already reached La-Nina according to the relative Nino index.

STATUS SUMMARY

1-month Rainfall status to October 2025

Base period: 1981-2021
Data source: MSWEP


Issued: 10/11/2025



3-month Rainfall status to October 2025

Base period: 1981-2021
Data source: MSWEP

Issued: 10/11/2025



6-month Rainfall status to October 2025

Base period: 1981-2021
Data source: MSWEP


Issued: 10/11/2025




12-month Rainfall status to October 2025

Base period: 1981-2021
Data source: MSWEP

Issued: 10/11/2025





Data source: MSWEP
 Method: Percentile
 © Commonwealth of Australia 2023, Australian Bureau of Meteorology, supported by COSPPac
 Model Run: 01/04/2023
 Base period: 1980-2021
 Shapefile data extracted from Flanders Marine Institute (2019), Maritime Boundaries Geodatabase: Maritime Boundaries and Exclusive Economic Zones (200NM), version 11. Available online at <http://www.marineregions.org/>.

Very dry conditions was recorded during the past 1, 3, 6, and 12 month timescale. Nauru experienced **below normal** rainfall and very dry conditions.

The maps above shows the past weather conditions from this month.

OUTLOOK SUMMERY

Chance of extreme rainfall for November 2025

Data source: ACCESS-S2
 Base period: 1981-2018

Model Run: 01/11/2025
 Issued: 05/11/2025



© Commonwealth of Australia 2025, Bureau of Meteorology, supported by COSPPac


Chance of extreme rainfall for November 2025 to January 2026

Data source: ACCESS-S2
 Base period: 1981-2018

Model Run: 01/11/2025
 Issued: 05/11/2025



© Commonwealth of Australia 2025, Bureau of Meteorology, supported by COSPPac



Data source: ACCESS-S2
 Issued: 03/05/2023
 © Commonwealth of Australia 2023, Australian Bureau of Meteorology, supported by COSPPac
 Model Run: 01/05/2023
 Base period: 1981-2018
 Shapefile data extracted from Flanders Marine Institute (2019), Maritime Boundaries Geodatabase: Maritime Boundaries and Exclusive Economic Zones (200NM), version 11. Available online at <http://www.marineregions.org/>.

Neutral conditions is expected for this month. Dry conditions is predicted to be easing off for this month.

For the next 3 month outlook from **November** to **January 2026**, **normal conditions** is also expected for **Nauru**.

The maps above shows chance of extreme rainfall for this month and the next 3 month outlook.

This in combination with the past status conditions suggests that ENSO is still in La-Nina conditions. However, La-Nina may form in the coming months across the tropical Pacific.

IMPACTS TABLE

After the specified period of below or above average rainfall, the following primary agricultural and hydrological variables and secondary socio-economic and health variables may to be impacted. Note the periods are estimates only. Allow for uncertainty associated with island size, topography, geology and soil type. Contact the relevant sector offices for further information on impacts.

VERY DRY TO SERIOUSLY DRY CONDITIONS (La Nina)

SECTOR	1 - month period most relevant for	3 - month period most relevant for	6 - month period most relevant for	12 - month period most relevant for
WATER	Small water tanks (5000ml), small creeks and streams (if present) Increased water demands	Small wells, Small streams (if present) Desalination plant unable to meet water demand	Boreholes, deeper wells, large water tanks.	Large water sources (e.g., rivers, artisan wells if present) Groundwater depletion
AGRICULTURE AND FOOD SECURITY	Shallow rooted crops (e.g., chillies, cherry tomatoes and eggplants)	Deeper rooted crops (e.g., bananas and guavas)	Small trees (including fruit trees e.g., breadfruit, pawpaw and sugarcane)	Large trees (including fruit trees e.g., coconut and breadfruit). Lime trees native trees Soil quality
HEALTH		Diarrhoea outbreak, asthma, skin rash.	Diarrhoea outbreak, asthma, skin rash.	Heat stroke, dehydration and conjunctivitis.
FISHERIES			Reduced tuna catches	
SOCIO- ECONOMIC			School closure	Increase household expenses on food.

VERY WET CONDITIONS (El Nino)

SECTOR	1 - month period most relevant for	3 - month period most relevant for	6 - month period most relevant for	12 - month period most relevant for
AGRICULTURE AND FOOD SECURITY	Reduced mangoes	Increased pig diseases due to poor shelter Soil quality (if shading nets poorly located)	Soil erosion Pandanus fruit rotting	
HEALTH	Increased incidences of dengue and gout	Increased incidences of dengue, flu, pneumonia and gout	Increased incidences of dengue, flu, pneumonia and gout	Increased incidences of dengue, flu, pneumonia and gout
FISHERIES	Increased tuna catches	coral bleaching		
SOCIO-ECONOMIC	Frequent road flooding Roof leaking Disrupts communication (increased cloud coverage) Wet laundry Mental health (stuck indoors)	Frequent flooding Disrupts phosphate mining Frequent electricity outage Increases and frequent shipping and flight delays (affect food supplies) School closures	Frequent flooding Disrupts phosphate mining Frequent electricity outage Increases and frequent shipping and flight delays (affect food supplies) School closures	Frequent flooding Disrupts phosphate mining Frequent electricity outage Increases and frequent shipping and flight delays (affect food supplies) School closures

RAINFALL MONITORING STATUS

The **rainfall status** maps are based on rainfall values from the MSWEP dataset which are then converted to the percentile index. The percentile index calculates the ranking of rainfall observed for a period against corresponding periods in the historical records for a particular timescale. MSWEP is a global precipitation product that combine rain gauges, satellite and ERA-5 reanalysis data and is provided at a 0.1° resolution. **Seriously dry** is defined as **Meteorological drought** assessed by rainfall

data only. It corresponds to rainfall for that period being in the bottom 10% of the historical record. **NO EXTREME** indicates that rainfall is within that middle 60% of historical observations for the respective timescale. In other words, rainfall that is **not extreme**. The 3 - 6 and 12 month timescale are more accurate representations of drought while the 1 - month timescale can be used to provide an indication of recent **dry (or wet)** spell conditions.

RAINFALL OUTLOOK

The **chance of extreme rainfall** maps are based on the likelihood of **very wet** or **very dry conditions**. This is equivalent to the chance that rainfall for that forecast period will be in the top or bottom 20% of historical observations for that selected period. The darker the shading, the more likely these extreme scenarios are. The white shading refers to a **low chance of extreme** which means the most likely scenario for that outlook period is for rainfall to be near **average** or **slightly below average** or **slightly above average** (not very wet nor very dry). The outlooks have been produced using the <http://www.bom.gov.au/climate/ahead/about/model/access.shtml>.

CONTACT FOR MORE INFO

Email us at nauru.met16@gmail.com for any concerns or suggestions

or contact the NMHS Director Mr. Graymea Ika at graymeaika1510@gmail.com (+674 558 8898)